

A 1+1 field theory spectrum from M theory

Maria Jose Rodriguez^{a*} and Pere Talavera^{b*}

^a Departament d'Estructura i Constituents de la Matèria,
Universitat de Barcelona, Diagonal 647, E-08028 Barcelona, Spain

^b Departament de Física i Enginyeria Nuclear,
Universitat Politècnica de Catalunya, Jordi Girona 1–3, E-08034 Barcelona, Spain

The spectrum of a 1+1 dimensional field theory with dynamical quarks is constructed. We focus in testing the possible brane embeddings that can support fundamental matter. The requirement on the wave function normalisation and the dependence on the quark mass of the quark condensate allow to discard most of the embeddings. We pay attention to some more general considerations comparing the behaviour of the non-compact theory at different dimensions. In particular we explored the possibility that the AdS/CFT duality “formalism” introduce a scale breaking parameter at $(1 + 1)d$ allowing the existence of *classical* glueballs and its possible relation with point-like string configurations. The screening effects and the appearance of a possible phase transition is also discussed.

August 2005

* majo@ecm.ub.es

* pere.talavera@upc.edu

1. Motivation and conclusions

There has been a lot of effort in trying to understand the low-energy dynamics of non-supersymmetric asymptotically free gauge theories from the supergravity duals (see for instance [1,2,3,4]). Most of these studies are focous at dimensions $d = 4, 3$ where confinement can only be understood as a dynamical feature while the case of dimension $d = 2$ is relegated to oblivion, probably because confinement is automatically incorporated in the theory even at perturbative level. The most salient point is precisely that the theory is fully resolvable in a suitable limit.

Field theory models at $d = 2$ while retaining some features of the four dimensional QCD theory as quark confinement and chiral symmetry breaking, still differs substantially in many crucial aspects from it: *i)* there are no dynamical gluons and hence strings are only build from matter quanta alone. *ii)* There is no chromomagnetic field, a key point to understand confinement in 3+1 dimensions and *iii)* it neither contains spin.

We want to elaborate here on the physical spectra, vectors, massive scalars and possible glueballs, one can find when dealing with one of these non-supersymmetric confining backgrounds in $d = 2$ [2]. We shall argue, that parallel to the field theory expectations, the supergravity model properties are as follows: there is spontaneous symmetry breaking, there are massless particles, the physical vectors acquires a mass. There are, obviously, still many lacking desired features but is encouraging to obtain some of the expected field theory results emerging from brane configurations.

In order to illustrate the field theory side we are after for, we review it briefly.

1.1. Field theory set up

The field theory parallel we want mainly to study is the 1+1 dimensions $SU(N_c)$ gauge theory coupled to matter field in the fundamental representation, [5,6] namely the 't Hooft model. Although our initial setup will be related to the $SU(N_c)$ gauge field coupled to static fermions in the adjoint representation [7].

The main difference between both settings are found in their interpretation as string models: while the 't Hooft model can be thought of as an open string model with a single rising Regge trajectory spectrum, the coupling to adjoint matter produces a kind of closed string with multiple Regge trajectories.

In the case of $SU(N_c)$ YM coupled to fundamental matter, our main purpose, the Minkowsky space action is taken to be

$$S = \int d^2x \text{Tr} \left[\bar{q} (i\gamma^\mu D_\mu - m_q) q - \frac{1}{2g_{\text{strong}}^2} F^{\mu\nu} F_{\mu\nu} \right], \quad (1.1)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + i[A_\mu, A_\nu]$ and the covariant derivative is $D_\mu \Phi = \partial_\mu \Phi + i[A_\mu, \Phi]$. In this case the quark-antiquark sector admits an infinite tower of confined color-singlets.

The reason for this solubility lies in the very defining features of the model. The large N_c limit eliminates all the sea quark contributions together with the non-planar gluon diagrams. On the other hand the fact of being in an axial gauge, $A_+ = 0$, and being the action independent of x_- derivatives allows to disentangle the gluon self-coupling by gauging away A_- . The only remaining Feynman diagrams in the 2-PI Green function are the rain-bow and ladder type, whose Schwinger-Dyson equation, giving the meson spectrum, can be solved in principle numerically.

The content of the paper is: we first start reviewing the construction of the non-supersymmetric background in a very synthetic way, it mainly should introduce the notation we shall follow and all the possible relevant embeddings. In sec. 3 a possible physical embedding providing matter in the fundamental is introduced. We check that from the initial two possibilities only one is feasible due to the normalisation at infinity. For the remaining solution we look for the relation between the chiral condensate and the quark mass. After comparison with the field theory and lattice calculations we find that the obtained results do not follow their trends. In accordance we turn to the evaluation of other embeddings in sec. 4. Once more the behaviour of the wave function at infinite imposes serious restrictions on the allowed solutions. For the remaining solutions we find the massive scalar and vector mesons spectra.

Thenceforth the subjects we treat has no special relation of been treating a 1+1 dimensional theory, but on the fact of having added fundamental matter. In sec. 5, following field theory arguments, we argue on the possible existence of classical glueballs. We present both, numerical and analytical evidence that the formalism can not account to their existence. Although we do not obtain glueball modes we check in sec. 6 the existence of oscillating modes on a “cigar” type configuration. This configuration was suggested to be related, at least in the four dimensional case, to the supergravity glueball spectra. The existence of the screening effect, hence the existence of at least two different phases, is discussed in sec. 7.

2. Non-extremal Dp-brane model

We shall introduce the main notation for the different quantities used in the text. The starting point is the metric and the dilaton field

$$ds^2 = h^{-1/2} dx_{||}^2 + h^{1/2} (dU^2 + U^2 d\Omega_{8-p}^2) , \quad e^\phi = (2\pi)^{2-p} g_{\text{YM}}^2 h^{(3-p)/4} . \quad (2.1)$$

The transverse space to the brane has dimension $9 - p$, and the wrap factor is given by

$$h^{-1} = \frac{U^{7-p}}{(g_{\text{YM}} \sqrt{d_p N_c})^2} . \quad (2.2)$$

In the remainder it will prove useful to define $R_p^{7-p} = (g_{\text{YM}} \sqrt{d_p N_c})^2$.

The non-extremal metric is obtained by imposing anti-periodic boundary conditions on the adjoint fermions, so that they become massive [2]

$$ds^2 = h^{-1/2} \left(dy_{\parallel}^2 + f(U) d\theta_2^2 \right) + h^{1/2} \left(\frac{dU^2}{f(U)} + U^2 d\Omega_{8-p}^2 \right), \quad (2.3)$$

where the function f

$$f(U) = 1 - \left(\frac{U_h}{U} \right)^{7-p}, \quad (2.4)$$

contains an IR scale, U_h , that breaks conformal invariance. Notice that at very high energy, $U \gg U_h$, (2.3) reduces to (2.1). To avoid the conical singularity the period of the compact θ_2 variable is chosen to be $\delta\theta_2 = 4\pi/(7-p)(R_p/U_h)^{(7-p)/2}U_h$. This compactification has a mass scale associated

$$m_{KK} = \frac{2\pi}{\delta\theta_2} = \frac{1}{2U_h} (7-p) \left(\frac{U_h}{R_p} \right)^{(7-p)/2}. \quad (2.5)$$

For energy scales lower than m_{KK} the theory is effectively p -dimensional. The transverse part of the Dp-brane can be parametrised by a set of coordinates $\vec{z} = (z^1, \dots, z^{9-p})$ suitable to write it as conformally flat

$$ds^2 = h^{-1/2} \left(dy_{\parallel}^2 + f(U) d\theta_2^2 \right) + K(\rho) d\vec{z} \cdot d\vec{z}, \quad (2.6)$$

where $\rho^2 = \vec{z} \cdot \vec{z}$ and $d\vec{z} \cdot d\vec{z} = d\rho^2 + \rho^2 d\Omega_{8-p}^2$. In order to obtain (2.6) one identifies

$$K(\rho) = h^{1/2} \left(\frac{U}{\rho} \right)^2, \quad (2.7)$$

and perform the change of variables $\partial_\rho U = U \sqrt{f(U)}/\rho$. When inserting (2.4) in the previous expression one gets

$$U(\rho) = \left[\frac{1}{2} \left(\frac{U_h \rho}{A} \right)^{\frac{7-p}{2}} + \frac{1}{2} \left(\frac{U_h A}{\rho} \right)^{\frac{7-p}{2}} \right]^{\frac{2}{7-p}}, \quad (2.8)$$

with A an integration constant. In order to conform the notation in [8] we set $(U_h/A)^{\frac{7-p}{2}} = 2$, from where

$$U(\rho) = \rho \left[1 + \frac{1}{4} \left(\frac{U_h}{\rho} \right)^{7-p} \right]^{\frac{2}{7-p}}. \quad (2.9)$$

To incorporate matter in the fundamental representation one embeds a probe-brane wrapping trivial circles on the transverse space of (2.6) [9]. Bearing that in mind we shall

depict the transverse space in a way that respect the symmetries of the would be embedded-brane. As previously mentioned we are interested in study the parallel of QCD_{1+1} . The responsible of the background will be a stack of D2-branes and the possible embeddings are given by the arrows

$$\begin{array}{llllllllll}
\text{D2:} & 0 & 1 & 2 & - & - & - & - & - & - \\
\text{D4:} & 0 & 1 & - & 3 & 4 & 5 & - & - & - \\
\text{D6:} & 0 & 1 & - & 3 & 4 & 5 & 6 & 7 & - \\
\text{D8:} & 0 & 1 & - & 3 & 4 & 5 & 6 & 7 & 8
\end{array}$$

Attending to the symmetries of the possible embedded Dm-probe brane the background field metric is written as

$$ds_{p=2}^2 = h^{-1/2} \left(ds^2 \left(E^{(1,1)} \right) + f(U) d\theta_2^2 \right) + K(\rho) (d\Omega_{m-1}^2 + d\Omega_{8-m}^2) , \quad (2.10)$$

Being from now on λ and r the radii of the transverse n-spheres to the Dp-brane, with $\rho^2 = \lambda^2 + r^2$.

3. Chiral symmetry breaking

The embedding of N_f Dm-probe brane in the ambient space of N_c Dp-brane ($p \leq m$) can be approximated by the DBI action

$$S_{\text{Dm}}^{(p)} = -\frac{1}{(2\pi)^m g_s \ell_s^{m+1}} \int d^{m+1} \sigma e^{-\phi} \sqrt{-\det g} , \quad (3.1)$$

if N_f is held fixed and $N_c \gg N_f$. In (3.1) g refers to the pullback metric. For later purposes the Dm-brane tension is defined as $T_{\text{Dm}} = ((2\pi)^m g_s \ell_s^{m+1})^{-1}$, and its world-volume coordinates will be parametrised by $\sigma^{0,\dots,m} = x^{0,\dots,p-1}, z^{p,\dots,m}$.

In the sequel, and to perform the embedding, we shall deal with an ansatz concerning the position of the Dm-brane in the space spanned for the latest n-sphere in (2.10). This n-sphere will be parametrised by the radii r and the angles ϕ^a ($a = 0, \dots, n-2$). Then the ansatz is chosen to be [10]

$$r(\lambda), \quad \phi^a = \text{fixed}, \quad \tau = \text{fixed}, \quad (3.2)$$

where the latest condition will be relaxed afterwards.

3.1. Embedding on D2-branes

The possible embeddings on a D2-brane bulk geometry were sketched previously and their action read

$$S_{\text{Dm}} = -T_{\text{Dm}} \int d^{m+1} \sigma \left(\frac{R_2}{\rho} \right)^{5(m-4)/4} \left(1 + \frac{U_h^5}{4\rho^5} \right)^{(16-m)/10} \lambda^{m-2} \sqrt{1 + \dot{r}^2} \sqrt{h_{\Omega_{m-2}}}, \quad (3.3)$$

with $h_{\Omega_{m-2}}$ the determinant on a unit $(m+1)$ dimensional sphere.

The classical equation of motion for the transverse, $r(\lambda)$, mode is

$$\begin{aligned} \frac{d}{d\lambda} \left[\frac{\lambda^{m-2}}{\rho^{5(m-4)/4}} \left(1 + \frac{1}{4} \left(\frac{U_h}{\rho} \right)^5 \right)^{(16-m)/10} \frac{\dot{r}}{\sqrt{1 + \dot{r}^2}} \right] &= -\frac{1}{16\rho^2} U_h^5 \\ \frac{\lambda^{m-2}}{\rho^{5m/4}} \left(1 + \frac{1}{4} \left(\frac{U_h}{\rho} \right)^5 \right)^{(6-m)/10} r \sqrt{1 + \dot{r}^2} &\left(3(4+m) + 20(m-4) \left(\frac{\rho}{U_h} \right)^5 \right). \end{aligned} \quad (3.4)$$

The BPS case is recovered by setting $U_h = 0$, then a particular solution to (3.4) is $r(\lambda) = \text{constant}$ reflecting that no force acts on the Dm-brane. To analyze further (3.4) is worth to rescale it as $\lambda \rightarrow U_h \lambda$, $r \rightarrow U_h r$, $\rho \rightarrow U_h \rho$. The differential equation is non-linear and it was not possible to find an analytic solution to (3.4) therefore we analyzed it asymptotically. More in concrete, we search for solutions in the asymptotic region with finite distance between the D2 and the Dm branes. That is $\lambda \rightarrow \infty$, $r(\lambda) \rightarrow r_\infty$. This in turn implies $\dot{r} \rightarrow 0$ and $\rho \sim \lambda$ which can be thought as the linearisation of the equation. With the above behaviour (3.4) becomes

$$\frac{d}{d\lambda} \left[\lambda^{3-m/4} \dot{r} \right] = -\frac{1}{16} r(\lambda) \frac{1}{\lambda^{m/4+4}} (3(4+m) + 20(m-4)\lambda^5). \quad (3.5)$$

Notice that $\lambda \rightarrow \infty$ appears as an irregular (regular) singular point in the D6-brane (D4-brane) embedding signaling some kind of illness in its solutions. For the two possible non-trivial embeddings one obtains

$$\text{D4:} \quad \frac{d}{d\lambda} [\lambda^2 \dot{r}] = -\frac{3}{2} \frac{r(\lambda)}{\lambda^5}, \quad \text{D6:} \quad \frac{d}{d\lambda} [\lambda^{3/2} \dot{r}] = -\frac{5}{2} \frac{r(\lambda)}{\lambda^{1/2}}, \quad (3.6)$$

while for the former one can find a normalisable solution for the radial coordinate

$$r(\lambda) = A \frac{1}{\sqrt{\lambda}} J_{-\frac{1}{5}} \left(\frac{\sqrt{6}}{5} \frac{1}{\lambda^{5/2}} \right) + B \frac{1}{\sqrt{\lambda}} J_{\frac{1}{5}} \left(\frac{\sqrt{6}}{5} \frac{1}{\lambda^{5/2}} \right), \quad (3.7)$$

the latter embedding presents an asymptotic oscillatory behaviour thus lacking any physical interpretation. We shall focus on the D4 case in the remainder.

The D4 embedding case reduces to, for arbitrary large λ values,

$$r(\lambda) = r_\infty + \frac{c}{\lambda}. \quad (3.8)$$

The physical interpretation of the coefficients r_∞ and c is the same as in the QCD_{3+1} case [8] : they are related to the quark mass and the chiral condensate respectively

$$m_q = \frac{U_h}{2\pi} r_\infty, \quad \frac{\delta \mathcal{E}}{\delta m_q} = \langle \bar{q}q \rangle = -8\pi^2 T U_h^2 c. \quad (3.9)$$

In order to gain more intuition on the solution, we inspect the region $r_\infty \gg$. This must be a small perturbation to the BPS state. Inserting $r(\lambda) = r_\infty + \delta r(\lambda)$ in (3.4) leads, at leading order in δr ,

$$\frac{d}{d\lambda} (\lambda^2 \delta \dot{r}) \approx -\frac{3}{2} r_\infty \frac{\lambda^2}{(r_\infty^2 + \lambda^2)^{7/2}}. \quad (3.10)$$

Integrating with the boundary conditions: *i)* $\dot{r}|_{\lambda=0} = 0$ and *ii)* $\delta r|_{\lambda \rightarrow \infty} = 0$ gives the final answer

$$r(\lambda) \approx r_\infty + \frac{3r_\infty^2 + 2\lambda^2}{10r_\infty^3 (r_\infty^2 + \lambda^2)^{3/2}}. \quad (3.11)$$

From the asymptotic result we can match the coefficients of the previous relation with those of (3.8) getting $c(r_\infty) \sim 1/(5r_\infty^3)$, i.e. the chiral condensate scales as $1/m_q^3$, for large quark masses. We shall comment on that behaviour latter on in connection to the field theory expectations. Notice that (3.11) has negative (positive) derivative for increasing positive (negative) values of λ , expecting then, a kind of bump or repulsion between the D4 and the D2-branes near the origin.




Fig. 1: In the l.h.s. panel there is plotted the D4 profile in the 8-9 plane. The D2-brane is the dashed line. In the r.h.s. we plotted the scaling of the quark condensate with respect to the quark mass.

In fig. 1 l.h.s. we depicted the profile of the D4 probe-brane in the 8 – 9 transverse plane. The D2-brane is located on the x-axis. In the r.h.s. we have evaluated numerically the coefficients c and r_∞ coming from (3.4). The numerical results corroborates all the previous analytical findings.

We can compare without further ado the previous findings with the corresponding field theory. At high energy or in processes involving heavy-quarks is well known that the use of summe-rules can provide the desired link between the hadron observables in terms of the fundamental degrees of freedom. For heavy-quarks, the quark condensate behaves as $\langle \bar{q}q \rangle \sim \langle FF \rangle / m_q^r$ been r a c-number fixed on dimensional grounds. The dimensional counting derived from (1.1) leads to: $[q] = E^{1/2}$ and $[F] = E$ hence it follows that $r = 1$, i.e. $\langle \bar{q}q \rangle \sim 1/m_q$. The previous expressions does not match with its purported supergravity dual.

4. $\overline{\text{Dm-D2-Dm}}$ system

It is clear that (3.2) is not able to reproduce the proper quark condensate in $1 + 1$ consequently we shall turn now to another ansatz involving the compact direction [8]. In the 4 dimensional case this was throughly explored in [11]. The embedding for the D4, D6 and D8 (generically denoted Dm) in the bulk of the D2 with $U(\theta_2)$ gives an action

$$S_{\text{Dm}} = -\frac{T_{\text{Dm}}}{g_{\text{YM}}^2} \int d^{m+1} \sigma \left(f(U) + \frac{h}{f(U)} (\partial_{\theta_2} U)^2 \right)^{1/2} U^{m-2} h^{(m-6)/4} \sqrt{h_{\Omega_{m-2}}}, \quad (4.1)$$

that does not depend explicitly on θ_2 . This last fact allows to obtain a first integral of motion

$$\theta_2(U) = \sqrt{f(U_0)U_0^{11-m/2}} \int_{U_0}^U \frac{dU}{h^{-1/2}f(U)\sqrt{f(U)U^{11-m/2} - f(U_0)U_0^{11-m/2}}}, \quad (4.2)$$

where U_0 is fixed by the condition $\partial_{\theta_2} U = 0$. From now on we choose without lost of generality $U_0 = U_h$.

For convenience we change variables. In the setup (4.1) the $U(\theta_2)$ coordinates are related to the cylindrical ones by

$$U^{7-p} = U_h^{7-p} + U_h^{5-p} r^2, \quad \theta = \frac{2\pi}{\delta\theta_2} \theta_2, \quad (4.3)$$

and these in turn with the Cartesian ones by the standard projection $y = r \cos \theta$, $z = r \sin \theta$. In the latter set, the BPS solution $U = U_h$ reads $y = 0$. In the following we shall perturb the metric around this BPS solution obtaining the spectra and checking its stability.

Instead of finding the quark mass dependence of the quark condensate as previously we turn to the direct evaluation of the spectra.

4.1. Worldvolume spectra

One of our aim is to obtain the low-energy spectrum in the afore presented theory, similarly to [12] a discrete pattern is expected. To work out the first order corrections to the spectra of the worldvolume fields it suffices to expand the action up to quadratic order. We shall deal with the correction to the scalar/pseudoscalar and gauge fields.

4.1.2. Vector mesons

For gauge fields the relevant part of the Lagrangian density can be written as

$$\mathcal{L} = -\tilde{T}_1 \alpha'^2 U^{-(14+m)/4} \left(2R_2^5 F_{\mu\nu} F^{\mu\nu} + 25 \frac{U^8}{U_h^3} F_{\mu z} F_{\nu z} \eta^{\mu\nu} \right). \quad (4.4)$$

Like we look for the single states in the S^{m-2} we have set to zero the component of the gauge in the compact space. This in turn eliminates any contribution of the Chern-Simon terms.

In order to find the spectrum we expand the field content of (4.4) in a complete set of eigenfunctions in the transverse coordinate z : $A_\mu(x^\mu, z) = \sum_n B_\mu^{(n)}(x^\mu) \psi_n(z)$, $A_z(x^\mu, z) = \sum_n \varphi_\mu^{(n)}(x^\mu) \phi_n(z)$. To normalise canonically the kinetic terms of the Yang-Mills (A_μ), the vectors fields (B_μ) and scalar or pseudoscalar particle (φ) we use the rescaled variable $Z = z/U_h$. Defining the function $K(U) = (U/U_h)^{7-p} = 1 + Z^2$ the gauge part of the action derived from (4.4) is proportional to

$$S_{\text{Dm}} \sim - \int dZ d^2x K^{-(14+m)/20} \left[\frac{1}{4} F_{\mu\nu}^{(r)} F^{\mu\nu(s)} \psi_r \psi_s + \frac{1}{2} K^{8/5} m_{KK}^2 B_\mu^{(r)} B^{\mu(s)} \partial_Z \psi_r \partial_Z \psi_s \right], \quad (4.5)$$

with the proper normalisation condition for the ψ and ϕ modes. The Born level e.o.m. for the ψ mode can be derived from (4.5)

$$-K^{(14+m)/20} \partial_Z \left(K^{(18-m)/20} \partial_Z \psi_r \right) = \lambda_r \psi_r, \quad (4.6)$$

where we have identify the squared mass of ψ with the eigenvalue λ .

With the use of (4.6) and the identification $\phi_n \rightarrow m_{kk}^{-1} / \sqrt{\lambda_n} \partial_Z \psi_n$ ($n \geq 1$), the modes with $n \geq 1$ can be gauged away by redefining $B_\mu^{(n)}$. This leads to the final action

$$S_{\text{Dm}} = - \int d^2x \left[\frac{1}{2} \partial_\mu \varphi^{(0)} \partial^\mu \varphi^{(0)} + \sum_{n \geq 1} \left(\frac{1}{4} F_{\mu\nu}^{(n)} F^{\mu\nu(n)} + \frac{1}{2} M_n^2 B_\mu^{(n)} B^{\mu(n)} \right) \right]. \quad (4.7)$$

The above expression indicates that the scalar/pseudoscalar and the Yang-Mills fields are massless while the vector field B acquires a mass $M_n^2 = \lambda_n m_{KK}^2$ dictated by the eigenvalues in (4.6).

In the remainder of this section we shall determine the massive vector spectrum. For that purpose we focus in the eigenvalues of (4.6). We solve (4.6) asymptotically, at $Z \gg 1$. This solution together with the boundary conditions at the horizon gives an allowed discrete set of λ_n .

To search for the asymptotic form of the wave function in (4.6) we apply the Frobenius method. The point $z \rightarrow \infty$ is a singular regular one. For $m = 4, 6$ we find two roots for the indicial equation. One of them corresponds to a non-normalisable solution while the other leads to $\psi \sim Z^{(m-8)/10}$. For the case $m = 8$ the roots of the indicial equation are degenerate and it does not contain normalisable wave functions. In the former case the eigenfunctions can be expanded in power series $\psi = \sum \alpha_k Z^{-2k/5 - (8-m)/10}$ with coefficients following from (4.6). Once the arbitrariness in the first one is removed by choosing $\alpha_0 = 1$, the asymptotic behaviour of ψ in (4.6) is fixed. The first coefficients are: $\alpha_1 = \alpha_2 = \alpha_4 = \alpha_7 = 0$, $\alpha_3 = 25 \lambda_n / (3m - 60)$, $\alpha_5 = (144 - 26m + m^2) / (20m - 560)$, $\alpha_6 = 625 \lambda_n^2 / (18(640 - 52m + m^2))$.

In addition to the asymptotic behaviour the regularity at the origin, $Z = 0$, demands ψ to be an even or an odd function. These constraints pin down a discrete set of eigenvalues in (4.6) corresponding to the vector masses depicted in table 2.

Eigenvalue	$m = 4$	$m = 6$
λ_1	0.309	0.149
λ_2	1.591	1.290
λ_3	3.811	3.349
λ_4	6.983	6.361

Table 2: The vector, axial-vector spectrum for different dimensional embeddings as obtained by (4.6).

Notice that in increasing the number of transverse directions the mass of the vector, axial-vector decreases in accordance with the absorption probabilities for emission on the brane.

The squared masses of the vectors and vector-axial fields are represented in fig. 2. For both quadratic expressions in terms of the principal quantum number, $a + bn + cn^2$ holds.

4.1.1. Massive scalar mesons

We are now in a position to obtain in a similar manner the massive scalar spectrum. For that purpose we excite the scalar sector with the ansatz $y(x^\mu, z)$. Notice that contrary to the assignment in [8] y does not depend on the S^{m-2} coordinates.




Fig. 2: vector and axial-vector squared masses as a function of the state. To guide the eye we have link the states with lines. Even if in appearance seems to appears a linear behaviour this is no the case.

In terms of the function K the induced Dm action takes the form

$$S_{\text{Dm}} = -\tilde{T} \int dZ d^2x \left(\frac{1}{2} K^{-(14+m)/20} \partial_\mu y \partial^\mu y + \frac{1}{2} m_{KK}^2 K^{(18-m)/20} \partial_Z y \partial_Z y + \frac{3}{5} m_{KK}^2 y^2 \right), \quad (4.8)$$

whose corresponding energy density is positive, ensuring the brane stability under small perturbations. To obtain a two-dimensional field theory description, we expand the scalar field in a complete set of eigenfunctions in the transverse direction: $y(x^\mu, z) = \sum_n \mathcal{U}^{(n)}(x^\mu) \rho_n(z)$. This allows to fix the equation of motion (e.o.m.) at tree level for the ρ mode

$$-K^{(14+m)/20} \left[\partial_Z \left(K^{(18-m)/20} \partial_Z \rho_n \right) - \delta \rho_n \right] = \lambda_n \rho_n, \quad \delta = \frac{6}{5}. \quad (4.9)$$

Notice that the previous expression is identical to (4.6) with the only exception of the second term in the squared brackets. Thus *naively taking the limit $\delta \rightarrow 0$ we shall recover the vector spectrum.*

All these ingredients together with the requirement of a canonical normalisation of the kinetic term fixes the lagragian density

$$\mathcal{L} = -\frac{1}{2} \sum_n \left(\partial_\mu \mathcal{U}^{(n)} \partial^\mu \mathcal{U}^{(n)} + M_n^2 \mathcal{U}^{(n)} \mathcal{U}^{(n)} \right), \quad M_n^2 = m_{KK}^2 \lambda_n. \quad (4.10)$$

In order to analyse the asymptotic behaviour for the different embeddings it is convenient to perform the change of variables $Z \rightarrow \omega^{-1/a}$, with a been an arbitrary positive c-number fixed only by the requirement of canceling possible poles at $\omega \rightarrow 0$. Then (4.9) becomes

$$\partial_\omega^2 \rho(\omega) + p(\omega) \partial_\omega \rho(\omega) + q(\omega) \rho(\omega) = 0, \quad (4.11)$$

with

$$\lim_{\omega \rightarrow 0} p(\omega) \rightarrow \frac{(m + 10a - 8)}{10 a \omega}, \quad \lim_{\omega \rightarrow 0} q(\omega) \rightarrow -\frac{1}{a^2} \left(\delta \omega^{-(20+(2+m)/a)/10} - \lambda \omega^{-2+6/(5a)} \right), \quad (4.12)$$

where we have only displayed the divergent terms. Notice that $\lim_{\omega \rightarrow 0} \omega p(\omega) \rightarrow 0$ while $\lim_{\omega \rightarrow 0} \omega^2 q(\omega)$ depends on the actual value of a . We can choose the coefficient a to cancel either the first or the second pole in $q(\omega)$ but not both simultaneously. In the vector case a was precisely chosen to make the λ term in $q(\omega)$ analytic. Now is not possible, provided $\delta \neq 0$. The conclusions are far obvious: the massive scalar wave function has not power series representation in the asymptotic region and hence it should be a non-analytic function. This implies that the solution of (4.9) does not go smoothly to the one obtained in (4.6) as $\delta \rightarrow 0$. To assess the correctness of this assertion we assume δ to be a free, small parameter and perform perturbation theory around the vector solution (4.6). If correct the previous findings, we expect to obtain the failure on the perturbation theory assumptions at some point. We write (4.11) as a Schrödinger like equation at zero energy with the potentials

$$V(x) + \delta v_{\text{pert}}(x) = \frac{(m - 18) ((2 + m)x^2 - 20)}{400(1 + x^2)^2} - \frac{\lambda}{(1 + x^2)^{8/5}} + \delta(1 + x^2)^{(m-18)/20}. \quad (4.13)$$

The correction to the ground state energy ($E = E_0 + \delta E_1$) reads

$$E_1 = \frac{\int_0^\infty dx \psi_0^2 (1 + x^2)^{(m-18)/10}}{\int_0^\infty dx \psi_0^2}, \quad (4.14)$$

where ψ_0 stands for the ground state (i.e. λ_1) wave function in (4.6).

The failure of perturbation theory can be traced back in the correction to the wave function ($\rho = \psi_0 + \delta \psi_1$)

$$\psi_1(x) = \psi_0(x) \int_0^x dz \frac{1}{(1 + z^2)^{(m-18)/20} \psi_0(z)^2} \int_0^z dt \psi_0(t)^2 (1 + t^2)^{(m-18)/10}. \quad (4.15)$$

The behaviour of ψ_1 is unbounded at large distance, contrary to that of ψ_0 . This contradicts the assumption, $\lim_{x \rightarrow \infty} \psi_1 \rightarrow 0$ where (4.14), (4.15) are built in. In conclusion the solution of (4.9) is not an analytic function of δ at the boundary.

To elaborate more on the physical relevance of the solution to (4.9) we apply the matching procedure [13]. For large, but finite, values of λ the solutions to (4.9) are not normalisable and hence the model does not contain nor scalars neither pseudoscalar massive particles. The fact that pseudoscalars particles are massless is not in conflict with been in a chiral symmetry broken phase [14]: the only necessary and sufficient criterion of

spontaneous chiral symmetry breaking (SB χ S) is a non-zero value of the left-right correlation function $i \lim_{m \rightarrow 0} \int d^4x \langle \Omega | T L_\mu^i(x) R_\nu^j(0) | \Omega \rangle = -\frac{1}{4} \eta_{\mu\nu} \delta^{ij} F_0^2$ been L_μ and R_μ the Noether currents generating the left and right chiral rotations respectively and F_0 , an order parameter, given by $\langle 0 | A_\mu^i | \pi^j \vec{p} \rangle = \delta^{ij} F_0 p_\mu$. There are certainly many other order parameters, such as local quark condensates $\langle \bar{q}q \rangle$, $\langle \bar{q} \sigma_{\mu\nu} F^{\mu\nu} q \rangle \dots$. A non-zero value of each of them by itself implies SB χ S, but the converse is not true i.e. SB χ S can take place ($F_0 \neq 0$) even if some of those condensate vanishes. In particular there is no available proof that $\langle \bar{q}q \rangle \neq 0$ is a necessary consequence of SB χ S.

In conclusion the spectrum of this second type of embedding reduces to massless pseudoscalars and massive vectors and axials. To wit its reliability we compare with the original 't Hooft model. There the full 2PI Green function amounts to

$$M_n^2 \varphi_n(x) = \left(\frac{m_q^2 - \beta^2}{x} + \frac{m_{\bar{q}}^2 - \beta^2}{1-x} \right) \varphi_n(x) - \beta^2 \int_0^1 dy \varphi_n(y) P \left[\frac{1}{(y-x)^2} \right], \quad (4.16)$$

with $x \in [0, 1]$ the momentum fraction carried by the quark in the light-cone coordinates and $\beta^2 = g_{\text{strong}}^2 / (2\pi)(N_c - 1/N_c)$. Furthermore β plays the analogous role in $(1+1)d$ of Λ_{QCD} in $(3+1)d$ [15]. The spectrum of (4.16) contains a single increasing Regge trajectory $M_\pi^2 \sim \beta^2 n$, ($n = 1, 2, \dots$), which is consistently found in the weak coupling regime, $m_q \gg g_{\text{strong}} \sim \frac{1}{\sqrt{N_c}}$ [16] and agrees with lattice calculations [17]. Then the lack of massive pseudoscalars in these type of embedding discards any possibility of these been a realistic duals to the $1+1$ 't Hooft model.

5. Supergravity glueballs

It was known long ago that classical Yang-Mills theory does not posses in $d = 2, 4$ finite energy non-singular time-independent solutions [18,19]. It was also proven that neither periodic in time solutions can exist [20]. It is then remarkable that in the AdS/CFT correspondence a *classical* background metric can account for these quantum objects at $d = 3, 4$ [21].

In $1+1$ the situation is as follows: in 1-dimensional space there is no possibility of having transverse gluons. The role of these transverse gluons in the glueball formation is to fix some scale that breaks the conformal invariance of the pure Yang-Mills theory. In the absence of this scale glueballs can not exist. It is not shocking then that models with adjoint fermions [22], higgs fields [23] or gravity [24] in $1+1$ -dimensions contains glueballs, provided these fields couple to gluons and gives the desired scale. The same role can be played by the use of a formalism incorporating some modification of the standard Yang-Mills density lagrangian that explicitly breaks scale invariance. This is the case of

the DBI action [25]. This was the precise case in the field theory side [26]. We shall check whether the model (2.3) reflects this peculiarity.

Following [2] we assume that the dilaton field couples to the $\text{Tr} F^2$ operator. In Einstein frame the e.o.m. for the scalar fluctuations are obtained via $\partial_\mu \left(\sqrt{g} g^{\mu\nu} \partial_\nu \tilde{\phi} \right) = 0$. In order to solve the previous expression and calculate the mass spectrum we follow an analogue procedure to the higher dimensional case [21] and choose a plane wave ansatz along the $\mathbb{R}^{(1,1)}$ directions, $\tilde{\phi} = e^{ik \cdot x} \chi(U)$. Using (2.3) the e.o.m. for the $d = 2$ massless dilaton boils down to

$$\partial_U \left(U^{-3/2} (U^5 - U_h^5) \chi'(U) \right) + U^{-3/2} M^2 R^5 \chi(U) = 0, \quad M^2 = -(k_1^2 - k_2^2). \quad (5.1)$$

The asymptotic solution depicts a normalisable term $\chi(U) \sim U^{-5/4} J_{5/6}(U^{-3/2}) \sim U^{-5/2}$. For large values of U (5.1) can be solved by a series solution with negative power $\chi = \sum_{n=0}^{\infty} a_n U^{-(n+5/2)} U_h^{-n}$. The first few non-vanishing coefficients are given by: $a_3 = -2M^2 R^5 / (33U_h^3)$, $a_5 = 1/3$, $a_6 = 2M^4 R^{10} / (1683U_h^6)$, where the arbitrary normalisation is fixed by the choice $a_0 = 1$.

The geometry must be regular at the horizon, thus we demand $\psi(U_h)$ to be regular. This fixes the possible values of M . We have scanned a wide range of possible values of M and found no solution satisfying the proper boundary conditions. In order to clarify the result we turn to a semi-classical evaluation of the spectra.

Semi-classical WKB approximation: A possible way to tackle (5.1) is to focus in highly excited states. There the depth of the potential is sufficient large for a semi-classical approximation be justified. Following [27] we perform first a functional change after which the potential can be written as a Schrödinger type equation at zero energy. Second we change variable to $U = F(\omega)$. The latter step forces to choose a new multiplicative factor to bring once more the remaining expression to a Schrödinger type one. All in all, applying the generic change of variables $\omega = \log(U^{7-p} - U_h^{7-p})$ the new potential reads (for $p = 2$)

$$V(\omega) = \frac{e^\omega (4U_h^5 + e^\omega)}{16(U_h^5 + e^\omega)^2} - \frac{M^2 R^5}{25} \frac{e^\omega}{(U_h^5 + e^\omega)^{8/5}}, \quad (5.2)$$

that is depicted as the full line in fig. 3 in comparison with the corresponding potentials at $d = 4, 3$. As one can see increasing the number of transverse coordinates the depth of the well is reduced. The question is whether for $d = 2$ still there is room to support at least the lowest state.

The potential in (5.2) takes the following asymptotic forms

$$V(\omega \ll) = \left(\frac{1}{4U_h^5} - \frac{M^2 R^5}{25U_h^8} \right) e^\omega + \left(-\frac{7}{16U_h^{10}} + \frac{8M^2 R^5}{125U_h^{13}} \right) e^{2\omega} + \dots \quad (5.3)$$




Fig. 3: The potential (5.2) . Dashed lines correspond to $d = 4, 3$ (short and long dashed respectively) while full line corresponds to $d = 2$. We have applied to each case the change of variables $\omega = \log(U^{7-p} - U_h^{7-p})$.

$$V(\omega \gg) = \frac{1}{16} - \frac{M^2 R^5}{25} e^{-3\omega/5} + \frac{U_h^5}{8} e^{-\omega} + \dots \quad (5.4)$$

Hence the classical turning point at large ω is approximately

$$\omega_+ \approx \frac{5}{3} \log \left(\frac{16M^2 R^5}{25} \right) , \quad (5.5)$$

while the inner turning point is located at $\omega_- \rightarrow -\infty$. In terms of the original variable U reads $\omega_- = R$.

Then expanding at leading order in $\frac{1}{M^2 R^5}$ the Borh-Sommerfeld expression leads to

$$\left(n - \frac{1}{2} \right) \pi \approx \frac{1}{5} M R^{5/2} \int_{-\infty}^{\omega_+} d\omega \frac{e^{\omega/2}}{e^{\omega} + U_h^5} + \mathcal{O} \left(\frac{1}{M^2 R^5} \right) . \quad (5.6)$$

A few remarks are in order: we discard the procedure in [27] that sets $\omega_+ \rightarrow \infty$, and subtract the corresponding piece as a perturbation because (5.6) becomes divergent. This is why we should bear in mind that the upper integration limit depends on the expansion parameter. The integration can be done analytically in terms of hypergeometric functions, but its results is not very illuminating. After this we are lead with an equality that for a given n can not be saturated for any value of M^2 .

Hence even if apparently the potential in fig. 3 has a minimum, it is not sufficient deep to hold even the ground state. We think that this does not enter in contradiction with the results of [26]. First of all because in deriving (5.1) one never makes use of the DBI action and secondly because the duality must give pure YM theory and not gluons coupled to gravity, hence the gravitons can not play the role of breaking any scale. Although this line

of reasoning, is striking that the role of the parameter U_h or the coupling of the dilaton to the Liouville field in the gravity side does not provide the desired breaking, probably this is an indication that classical glueballs are protected in two dimensions also in the supergravity side.

6. Breathing modes: pulsating string on $\mathbb{R}^+ \times \mathbb{R}^+$

In the following we shall inspect, for the general setup (2.3), a point-like string configuration that corresponds to a particle moving on the meridian of a 2-dimensional “cigar” shape surface, $U(\tau), t(\tau)$ with the rest of coordinates constant. In $3+1$ dimensions it was shown that this configuration was directly linked with the glueball spectra at leading order [28]. In our setup glueballs do not exist, but as we shall see a pulsating configuration still exists. In the conformal gauge the solutions to the e.o.m. and the Virasoro constraints describe null geodesics of (2.3) with solutions dictated by

$$\dot{t} = \frac{c}{2} \left(\frac{R_2}{U} \right)^{(7-p)/2}, \quad \dot{U} = \pm \frac{c}{2} \sqrt{f(U)}. \quad (6.1)$$

This corresponds to an harmonic oscillation with target space period

$$\Delta T = 2 \int_{U_h}^{\infty} dU \left(\frac{R_p}{U} \right)^{(7-p)/2} \frac{1}{\sqrt{f(U)}}, \quad (6.2)$$

and angular frequency $\omega_0 = 2\pi/\Delta T$.

To determine the spectrum we work, for convenience, with the Nambu-Goto action considering a general setting $U(\tau), t(\tau), \theta_2 = m\sigma$. With this ansatz the lagrangian density for (2.3) reduces to $\mathcal{L} = -m(h^{-1}f(U) - \partial_\tau U^2)^{1/2}$, that with the change of variables

$$\frac{d\zeta}{dU} = \frac{R^{(7-p)/2}}{\sqrt{U^{7-p} - U_h^{7-p}}}$$

becomes $\mathcal{L} = -m\sqrt{h^{-1}f(\zeta)}(1 - \partial_\tau \zeta^2)$. Its associated hamiltonian defines a one dimensional system with potential $V = m^2 h^{-1/2} f(\zeta)/2$. The semi-classical energy levels are given by

$$\begin{aligned} \left(2n + \frac{1}{2}\right) &\approx \int d\zeta \sqrt{E_n^2 - m^2 h^{-1/2} f(\zeta)} \\ &= 2 \int_{U_h}^{U_1(m)} \left(\frac{R}{U} \right)^{(7-p)/2} \sqrt{E_n^2 f(U)^{-1} - m^2 \left(\frac{U}{R} \right)^{(7-p)/2}}, \end{aligned} \quad (6.3)$$

where the upper turning point, $U_1(m)^{7-p} = (E^2/m^2)R^{(7-p)/2} - U_h^{7-p}$, diverges in the massless limit. In this case and for large energy states (6.3) reduces to $E_n \approx n\omega_0$ that for $p = 2$ gives

$$E_n \approx -\sqrt{\pi}n \left(\frac{U_h^3}{R^5} \right)^{1/2} \frac{\Gamma(-1/5)}{\Gamma(3/10)}. \quad (6.4)$$

Despite its appearance is positive defined.

7. Quark-antiquark potential

The Wilson loop is in YM the order parameter signalling confinement. In the case of adding matter it still can be used for this purpose. The case of YM theory coupled to adjoint matter was extensively studied in [29]. For the case at hand, addition of fundamental matter supported on probe-branes, the expressions for the energy and quark-antiquark separation are identical to those depicted in [29] with the only exception that an upper cutoff (given by the brane position in the transverse space) appears in the integration [30]. For finite transverse distance we have

$$E_{\text{unsub}} = \frac{U_0}{\pi} \int_1^{U_{\text{Dm}}/U_0} dy \frac{y^{7-p}}{\sqrt{(y^{7-p}-1)(y^{7-p}-\lambda)}}, \quad (7.1)$$

$$L = 2 \frac{R_p^{(7-p)/2}}{U_0^{(5-p)/2}} \int_1^{U_{\text{Dm}}/U_0} dy \frac{1}{\sqrt{(y^{7-p}-1)(y^{7-p}-\lambda)}}, \quad (7.2)$$

with $\lambda = (U_h/U_0)^{7-p}$. In the case of dealing with probe branes at infinity one has to subtract the bare quark mass in order to obtain a finite result

$$E_{\text{sub}} = E_{\text{unsub}} - \frac{U_0}{\pi} \int_1^{U_{\text{Dm}}/U_0} dy \sqrt{\frac{y^{7-p}}{y^{7-p}-\lambda}}. \quad (7.3)$$

We turn first to the asymptotic values. The long and short string limits can be attained when the turning point approaches the horizon, $\lambda \rightarrow 1^+$ and $U_0 \rightarrow U_{\text{Dm}}^-$ respectively. At leading order

$$E_{\text{long}} \approx \frac{1}{2\pi} \left(\frac{U_h}{R_p} \right)^{(7-p)/2} L + \dots, \quad E_{\text{short}} \approx \frac{1}{2\pi} \left(\frac{U_{\text{Dm}}}{R_p} \right)^{(7-p)/2} L + \dots \quad (7.4)$$

Thus there is an universal linear behaviour at both energy ends. We stress that, is suggestive, the interpretation of this phenomena at high-energy as a color screening effect: while in pure YM at large- N_c deconfinement by color screening is suppressed, [31], one can argue that is the addition of fundamental matter the responsible of this shape [32]. To substantiate furthermore this claim we increasing the distance between the probe brane with the

original stack. Then this linear behaviour disappears recovering the usual coulomb potential[†]. Eventhough we must bear in mind that we have only deal with the bosonic sector and the fermionic corrections are unavoidable for a correct treatment. Notice also that both expressions in (7.4) are afflicted with some drawbacks: increasing indiscriminately the energy we probe the compact dimension in S^1 , while in the other extreme one can neither trust all the way to the infrared limit. In decreasing the energy at the point where $U \sim g_{\text{YM}}^2$ the system description is in terms of $M2$ branes [33].

As second and concluding remark it is instructive to look at the concavity conditions of the inter-quark potential [34]

$$\frac{dE_{\bar{q}q}}{dL} > 0, \quad \frac{d^2 E_{\bar{q}q}}{dL^2} \leq 0. \quad (7.5)$$

These have already been study in several systems [35]. Contrary to the systems studied in the above reference our E vs. L relations show a smooth behaviour. As one can verify $\partial_{U_0} L$ is a monotonic decreasing function of U_0 with zeros lying outside the physical region.




Fig. 4: Energy as a function of the string turning point U_0 as (7.1) and (7.3). Short (long) dashed curve corresponds to $p = 4, (3)$, while the full curve to $p = 2$. We have set $R = U_h = 1$ for illustrative purposes.

In fig. 4 we depict the energy versus the string turning point using a subtracted and unsubtracted relation. Notice that the expression (7.3) has an intermediate maximum while (7.1) only takes its maximum in one of the edges. Like the difference between the two procedures is just a renormalisation scheme choice both must be equivalent, hence we conclude that with the addition of fundamental matter the shape of the subtracted energy can not be interpreted as a phase transition phenomena. These values of the energy must be

[†] One needs to subtract then the corresponding bare quantities.

compared with the ones needed for the opening of the compact dimensions, corresponding roughly to energies of the order of the the Kaluza-Klein masses (2.5) $m_{KK} = 1.5, 2, 2.5$ for $p = 4, 3, 2$ respectively. While in (7.1) these suppose a constraint for the allowed region in U_0 , only short strings are allowed, seems that (7.3) includes also the long string case.

The expression of the energy as a function of the inter-quark separation for (7.1) can be easily worked out

$$\frac{dE_{\text{unsub}}}{dL} = \frac{\lambda}{2\pi} U_0^{(7-p)/2} + \left(\frac{E_{\text{unsub}}}{U_0} - \frac{\lambda}{2\pi} \frac{U_0^{(5-p)/2}}{R_p^{(7-p)/2}} L \right) \frac{1}{\partial_{U_0} L}. \quad (7.6)$$

8. Summary

We have inspected the matter content of a 1+1 dimensional field theory with the hope of finding some resemblance with QCD_{1+1} that we know to be fully solvable in the strong coupling regime. From the two possible general supergravity ansatz we have studied the first one does not lead to the correct relation between the quark condensate and the quark mass dependence. While for the second type of embedding the spectrum reduces to massless pseudoscalars and massive vectors and axials.

Even if at first sight none of the trials to embed matter in the fundamental representation seems successful, the model contains several interesting properties as confinement and chiral symmetry breaking. Probably the lesson to learn is that the most sensible way to obtain a reliable model of QCD_{1+1} is not to directly compactify on an S^6 but compactify QCD_{3+1} or QCD_{2+1} on an S^2 or on an S^2 respectively. This would lead hopefully to the proper relation between the quark condensate and quark mass in the first type of embeddings discussed.

Together with this, we have looked for classical realisation of glueball in the D2-brane. Notice that this scalar sector is not the same as the one obtained in sec. 4.1.1. The former corresponds to the Kaluza-Klein singlets states over the initial S^6 , while the latter is obtained by exciting modes on the probe brane. These scalars are protected to appear in accordance to the field theory expectations. We have compared the difference with the same *family* of models but with different compact dimensions and elucidated, based on a semi-classical analysis, the reasons that leads to the lack of this kind of scalars.

Furthermore we have checked that even if glueballs are absent one can obtain a cigar-like configuration. In $d = 4$ the spectrum of this configuration reduces to that of the glueballs. This seems to be an exceptional case and we do not find deeper explanation for the coincidence.

We have also looked at the quark-antiquark potential. The behaviour, in particular the extrema, of the energy as a function of the string turning point depends on a point dependent finite subtraction. The difference between both procedures, subtracted and




Fig. 5: The full (dashed) line describes the subtracted (unsubtracted) energy as a function of the string length.

unsubtracted, is a renormalisation scheme choice, thus we conclude that the shapes in fig. 5 can not be interpreted as a phase transition.

Acknowledgements

We thank J. Boronat, J. Casulleras, T. Sakai, J. Soto, S. Sugimoto and O. Varela for discussions. Also A. Cotrone for suggestions. This work is partially supported by the grants CYT FPA 2004-04582-C02-01, CIRIT GC 2001SGR-00065 and by the European Community's Human Potential Programme under contract MRTN-CT-2004-005104 'Constituents, fundamental forces and symmetries of the universe'. M. J. R. wants to thank the Generalitat of Catalunya for her FI-IQUC research grant.

References

- [1] D. J. Gross and H. Ooguri, “Aspects of large N gauge theory dynamics as seen by string theory,” *Phys. Rev. D* **58**, 106002 (1998) [arXiv:hep-th/9805129].
- [2] E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” *Adv. Theor. Math. Phys.* **2**, 505 (1998) [arXiv:hep-th/9803131].
- [3] J. Polchinski and M. J. Strassler, “The string dual of a confining four-dimensional gauge theory,” arXiv:hep-th/0003136.
- [4] O. Aharony, “The non-AdS/non-CFT correspondence, or three different paths to QCD,” arXiv:hep-th/0212193.
- [5] G. 't Hooft, “A Two-Dimensional Model For Mesons,” *Nucl. Phys. B* **75**, 461 (1974).
- [6] C. G. . Callan, N. Coote and D. J. Gross, “Two-Dimensional Yang-Mills Theory: A Model Of Quark Confinement,” *Phys. Rev. D* **13**, 1649 (1976).
- [7] S. Dalley and I. R. Klebanov, “String spectrum of (1+1)-dimensional large N QCD with adjoint matter,” *Phys. Rev. D* **47**, 2517 (1993) [arXiv:hep-th/9209049].
- [8] M. Kruczenski, D. Mateos, R. C. Myers and D. J. Winters, “Towards a holographic dual of large- $N(c)$ QCD,” *JHEP* **0405**, 041 (2004) [arXiv:hep-th/0311270].
- [9] A. Karch and E. Katz, “Adding flavor to AdS/CFT,” *JHEP* **0206**, 043 (2002) [arXiv:hep-th/0205236].
- [10] J. Babington, J. Erdmenger, N. J. Evans, Z. Guralnik and I. Kirsch, “Chiral symmetry breaking and pions in non-supersymmetric gauge / gravity duals,” *Phys. Rev. D* **69**, 066007 (2004) [arXiv:hep-th/0306018].
- [11] T. Sakai and S. Sugimoto, “Low energy hadron physics in holographic QCD,” arXiv:hep-th/0412141.
- [12] G. T. Horowitz and H. Ooguri, “Spectrum of large N gauge theory from supergravity,” *Phys. Rev. Lett.* **80**, 4116 (1998) [arXiv:hep-th/9802116].
- [13] C. M. Bender, S. A. Orszag, “Advanced Mathematical Methods for Scientists and Engineers,” Springer Verlag (1991)
- [14] J. Stern, “Two alternatives of spontaneous chiral symmetry breaking in QCD,” arXiv:hep-ph/9801282.
- [15] B. Grinstein and R. F. Lebed, “Explicit quark-hadron duality in heavy-light meson weak decays in the 't Hooft model,” *Phys. Rev. D* **57**, 1366 (1998) [arXiv:hep-ph/9708396].
- [16] A. R. Zhitnitsky, “On Chiral Symmetry Breaking In QCD In Two-Dimensions ($N(C) \rightarrow \text{Infinity}$),” *Phys. Lett. B* **165**, 405 (1985) [*Sov. J. Nucl. Phys.* **43**, 999.1986 YAFIA,43,1553 (1986 YAFIA,43,1553-1563.1986)].
- [17] R. Narayanan and H. Neuberger, “The quark mass dependence of the pion mass at infinite N ,” arXiv:hep-lat/0503033.

- [18] S. Deser, “Absence Of Static Solutions In Source - Free Yang-Mills Theory,” *Phys. Lett. B* **64**, 463 (1976).
- [19] H. Pagels, “Absence Of Periodic Solutions To Scale Invariant Classical Field Theories,” *Phys. Lett. B* **68**, 466 (1977).
- [20] S. R. Coleman, “There Are No Classical Glueballs,” *Commun. Math. Phys.* **55**, 113 (1977).
- [21] C. Csaki, H. Ooguri, Y. Oz and J. Terning, “Glueball mass spectrum from supergravity,” *JHEP* **9901**, 017 (1999) [arXiv:hep-th/9806021].
- [22] F. Antonuccio and S. Dalley, “Glueballs from (1+1)-dimensional gauge theories with transverse degrees of freedom,” *Nucl. Phys. B* **461**, 275 (1996) [arXiv:hep-ph/9506456].
- [23] R. F. Dashen, B. Hasslacher and A. Neveu, “Nonperturbative Methods And Extended Hadron Models In Field Theory. 3. Four-Dimensional Nonabelian Models,” *Phys. Rev. D* **10**, 4138 (1974).
- [24] R. Bartnik and J. McKinnon, “Particle - Like Solutions Of The Einstein Yang-Mills Equations,” *Phys. Rev. Lett.* **61**, 141 (1988).
- [25] A. A. Tseytlin, “On non-abelian generalisation of the Born-Infeld action in string theory,” *Nucl. Phys. B* **501**, 41 (1997) [arXiv:hep-th/9701125].
- [26] D. Gal'tsov and R. Kerner, “Classical glueballs in non-Abelian Born-Infeld theory,” *Phys. Rev. Lett.* **84**, 5955 (2000) [arXiv:hep-th/9910171].
- [27] J. A. Minahan, “Glueball mass spectra and other issues for supergravity duals of QCD models,” *JHEP* **9901**, 020 (1999) [arXiv:hep-th/9811156].
- [28] J. M. Pons, J. G. Russo and P. Talavera, “Semiclassical string spectrum in a string model dual to large N QCD,” *Nucl. Phys. B* **700**, 71 (2004) [arXiv:hep-th/0406266].
- [29] A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, “Wilson loops, confinement, and phase transitions in large N gauge theories from supergravity,” *JHEP* **9806**, 001 (1998) [arXiv:hep-th/9803263].
- [30] A. Paredes and P. Talavera, “Multiflavour excited mesons from the fifth dimension,” *Nucl. Phys. B* **713**, 438 (2005) [arXiv:hep-th/0412260].
- [31] J. Greensite and M. B. Halpern, “Suppression Of Color Screening At Large N,” *Phys. Rev. D* **27**, 2545 (1983).
- [32] D. J. Gross, I. R. Klebanov, A. V. Matytsin and A. V. Smilga, “Screening vs. Confinement in 1+1 Dimensions,” *Nucl. Phys. B* **461**, 109 (1996) [arXiv:hep-th/9511104].
- [33] N. Itzhaki, J. M. Maldacena, J. Sonnenschein and S. Yankielowicz, “Supergravity and the large N limit of theories with sixteen supercharges,” *Phys. Rev. D* **58**, 046004 (1998) [arXiv:hep-th/9802042].
- [34] C. Bachas, “Convexity Of The Quarkonium Potential,” *Phys. Rev. D* **33**, 2723 (1986).
- [35] A. Brandhuber and K. Sfetsos, “Wilson loops from multicentre and rotating branes, mass gaps and phase structure in gauge theories,” *Adv. Theor. Math. Phys.* **3**, 851 (1999) [arXiv:hep-th/9906201].